



Detox - IV Drip

What is Detox IV?

Natural detoxification is carried out primarily by the liver in conjunction with the kidneys and colon. The liver acts as a filtration system - it is the site for converting toxins into waste products, cleansing the blood, and metabolizing various nutrients and medications.

Toxic substances are conjugated into forms that can be safely excreted from the body in the stool and urine. However, toxic overload from pesticides, heavy metals, medications, alcohol, poor diet, and stress often overburden the liver, requiring additional support for proper detoxification.

The Detox Drip is formulated with vitamins and minerals specific to the liver to support its natural detoxification pathways.

Uses

- Support detoxification pathways
- Skin conditions
- Environmental toxin exposure
- Brain fog, memory impairment
- Prolonged metabolic stress

Side Effects

Common

- Bruising at IV access site
- Hypoglycemia
- Fatigue
- Warming sensation
- Headache
- Temporary worsening of symptoms
- Increased urination

Rare

- Allergic reaction
- Edema
- Chills, sweats
- Rash
- Diarrhea
- Nausea
- Exhaustion
- Flu-like symptoms
- Achiness
- Irritability

Ingredients

- Ascorbic Acid
- Calcium Chloride
- Magnesium Chloride
- Taurine
- Molybdenum
- M.I.C.
- Sodium Bicarbonate
- Dexpanethenol
- Pyridoxine
- B-100 Complex
- Methyl-B12
- Folinic Acid
- Selenium
- Zinc Sulfate
- Lymphdiaral
- Quassia
- Sterile Water

Procedure

This IV Drip is 2 parts – stacked together and done one after another through the same IV access site.

The first drip is high in vitamins, minerals, and amino acids. The second is a glutathione push which helps to neutralize toxins and free radicals by binding to them, making it easier for the body to detoxify, excrete, and decrease inflammation.

Duration

1.-1.5 hours

Detox - IV Drip



Patient Prep

Please arrive on time and well-hydrated. Please ensure that you have eaten prior to the intravenous treatment.

Please inform a lab technician and/or your doctor if you have any allergies; have sensitivity to sulphur containing foods; have a history of any reaction to intravenous treatment; are pregnant or breastfeeding; have kidney or liver disease; are taking blood sugar lowering medications, thyroid medications, or cancer medications; or have any concerns regarding this treatment.

Please consult with your ND if your condition persists or worsens.

If you have any questions, please contact us at (604) 738-1012, ext.1

Notes
